

## ON EVOLUTION OF STATIONARY PROCESSES NEAR THE ORIGINS OF EXCITATION

<sup>1</sup>Sergiyenko L.S., <sup>2</sup>Nesmeyanov A.A.

<sup>1</sup>National research Irkutsk state technical university, Irkutsk;

<sup>2</sup>East-siberian institute of Ministry of Internal Affairs of Russia, Irkutsk, e-mail: lusia\_ss@mail.ru

In three-dimensional Euclidean space we study elliptic system of four equations in quotient derivatives of the fourth order with four unknowns that parabolically degenerates on the coordinate axis. In certain classes of function smoothness an existence of the unique (with precision up to random constant) limited solution of the system that meet the fixed boundary terms in cylinder, symmetric to the line of degeneration, is proved. The proof is carried out through the reduction of the results of differentiation and integration of the system equations and dividing variables according to classic algorithm of Fourier. While calculating coefficients of the line that represent the solution of common linear differential equation of the second order with a special point in the centre of the research area that is a consequence of the studied system, special multinomials of so-called triangle form were built. The results can be used while modeling stationary processes in asymmetric solenoid speed field.

**Keywords:** stationary processes, excitation

Let the stationary process be presented as a linear operator equation

$$I \operatorname{rot} T + B \operatorname{grad} s = 0, \quad \operatorname{div} T = 0, \quad (1)$$

where components  $u, v, w$  of the vector  $T$ , and scalar function  $s$  are dependent variables of arguments  $x, y, z, I$  and  $B$  are given matrixes

$$I = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \quad B = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & f \end{bmatrix},$$

$$f = \frac{(x^2 + y^2)^l}{r^2 + (x^2 + y^2)^k}.$$

With positive values of parameters  $l, k, r$  [1–3, 6].

As we approach the origin of the excitation, the studied fixed process starts to alter its structure – elliptic system (1) parabolically degenerates in the multiplicity  $x^2 + y^2 = 0$ , and for it common classic setting of problems of modern mathematical physics become incorrect [7].

### The first boundary problem.

Let us study the behavior of the system (1)

$$u_x + v_y + w_z = 0, \quad w_z - u_x - s_y = 0,$$

$$v_z - u_y + f(x, y)s_z = 0, \quad s_x - v_z + w_y = 0.$$

Near the degeneration line  $x = y = 0$  that is contained in cylinder

$$D = \{(x, y): x^2 + y^2 < R^2, 0 < z < z_0\}$$

$$\Delta w = 0, \quad s_{xx} + s_{yy} + f(x, y) s_{zz} = 0,$$

$$u(x, y, z) = \int_0^z [w_x(x, y, \xi) - s_y(x, y, \xi)] d\xi + J(y, z);$$

$$u(x, y, z) = \int_0^z [w_y(x, y, \xi) + s_x(x, y, \xi)] d\xi + Q(y, z).$$

with side surface  $\Gamma$ , upper  $\Gamma_1$  and bottom  $\Gamma_0$  bases. Let us define  $D_0$  as a part of axis  $OZ$ , that lies in  $\bar{D}$ .

*Problem 1.*

Find all conditions of existence and uniqueness in area  $D$  of the limited on the multiplicity of degeneration  $D_0$  solution of the system (1).

While  $l > 0, k = r = 0$ , conditions of the correctness of the problem are defined in [4].

In the presented work under terms

$$v|_{\partial\Gamma_0} = h, \quad w|_{\partial D} = g,$$

$$s|_{\Gamma_0 \cup \Gamma_1} = 0, \quad s|_{\Gamma} = q, \quad (2)$$

where  $h \in C_{0+\alpha}(\partial\Gamma_0); g \in C_{1+\alpha}(\partial D);$

$q \in C^2(\bar{\Gamma}); q|_{\partial\Gamma_0} = 0.$

The following is proved.

**Theorem 1.** While  $l = k = 1, u, r = R$  in classes of the function smoothness

$$(u, v) \in C^1(D/D_0) \cap C(\bar{D}/D_0),$$

$$w \in C^2(D) \cap C(\bar{D}),$$

$$s \in C^2(D/D_0) \cap C(\bar{D}/D_0)$$

exists a limited near the degeneration axis solution  $(u, v, w, s)$  of the problem (1)–(2), where  $u$  and  $v$  are defined with a precision up to random constant summand, and  $u, w,$  and  $s$  – in a single way.

Through a reduction of the results of differentiation and integration of the equations (1) according to the corresponding variables we obtain a system:

Functions  $J(y, z)$  and  $Q(y, z)$  are defined precisely up to the random constant summand from the system

$$\begin{aligned} Q_y + J_x &= -\lim_{z \rightarrow 0} w_z; \\ Q_x - J_y &= f(x, y) \lim_{z \rightarrow 0} s_z. \end{aligned}$$

The main point in the proof of the theorem is:

**Lemma 1.** Boundary problem

$$\begin{aligned} s_{xx} + s_{yy} + \frac{x^2 + y^2}{R^2 + x^2 + y^2} s_{zz} &= 0; \quad (3) \\ s|_{\Gamma_0 \cup \Gamma_1} &= 0, \quad s|_{\Gamma} = q, \\ q \in C^2(\bar{\Gamma}), \quad q|_{\partial\Gamma} &= 0. \quad (4) \end{aligned}$$

Has a unique solution in the cylinder  $D$ , that is limited while  $(x^2 + y^2) \rightarrow 0$ .

The proof is carried out according to the classic algorithm of Fourier where at first variables  $z$  and  $(x, y)$  are divided. The solution is built as line

$$s(x, y, z) = \sum_{n=0}^{\infty} b_n(z) a_n(x, y). \quad (5)$$

We obtain

$$\begin{aligned} b_n'' + \lambda_n b_n &= 0; \\ \Delta a_n - \frac{x^2 + y^2}{R^2 + x^2 + y^2} a_n &= 0. \quad (6) \end{aligned}$$

$$\gamma_{4+2i} = (-1)^i \frac{\mu^2}{4(i+2)(i+2+m)} \left[ 1 + \sum_{j=0}^{\alpha_i-2} \left( \frac{\mu}{2} \right)^{\alpha_i-2j} P_{ij} \right], \quad (11)$$

where

$$\mu = \frac{\pi n}{4}; \quad \alpha_i = \frac{2i+3+(-1)^i}{2}; \quad P_{00} = P_{10} = 0, \quad (12)$$

and while  $i \geq 2$

$$P_{ij}(m) = \sum_{\tau_j=0}^{i-2-2j} \sum_{\tau_{j-1}=0}^{\tau_j} \dots \sum_{\tau_0=0}^{\tau_1} \prod_{\eta=0}^j \frac{1}{(2+2\eta+\tau_{j+1-\eta})(2+2\eta+\tau_{j+1-\eta})}. \quad (13)$$

The solution of the equation (3) in cylindrical coordinates will look as:

$$s(\rho, \varphi, z) = \sum_{m,n=0}^{\infty} \left\{ \frac{\Psi_{mn}(\rho)}{\Psi_{mn}(R)} \sin \frac{n\pi z}{z_0} [A_{mn} \cos(m\varphi) + B_{mn} \sin(m\varphi)] \right\}. \quad (14)$$

Degeneration into trigonometric line of the given function on the surface  $\bar{\Gamma}$  function

$$q(\varphi, z) = \sum_{m,n=0}^{\infty} \sin \frac{m\pi z}{z_0} [A_{mn} \cos(m\varphi) + B_{mn} \sin(m\varphi)]. \quad (15)$$

Let us define the values of Fourier coefficients:

$$A_{0m} = \frac{2}{\pi R z_0} \int_{\Gamma} q(\varphi, z) \sin \frac{m\pi z}{z_0} d\Gamma, \quad m = 0, 1, 2, \dots,$$

Under boundary terms  $b_n(0) = b_n(z_0) = 0$  и  $\lambda_n = n\pi z/z_0$ ,  $n = 1, 2, 3, \dots$ , from the first equation (6) we find

$$b_n(z) = \sin \frac{n\pi z}{z_0}. \quad (7)$$

Then, from the second equation (6) in polar coordinates  $(\varphi, \rho)$  we define a solution of the view

$$a_n = \Phi_n(\varphi) \cdot \Psi_n(\rho).$$

We obtain the system

$$\begin{aligned} \Phi_n'' + \gamma_n \Phi_n &= 0; \\ \rho^2 \Psi_n'' + \rho \Psi_n' - \left( \frac{\rho^4}{R^2 + \rho^2} + \gamma_n \right) \Psi_n &= 0. \quad (8) \end{aligned}$$

From the first equation (8) while  $\gamma_n = m^2$ ,  $m = 0, 1, 2, \dots$  single periodic solution in shape of harmonic superpositions

$$\Phi_{nm}(\varphi) = A_{nm} \cos(m\varphi) + B_{nm} \sin(m\varphi). \quad (9)$$

With each focused  $n$  the second equation of the system (8) always has an integral as a line

$$\Psi_{mn}(\rho) = \rho^m \left( 1 + \sum_{i=0}^{\infty} \gamma_{4+2i} \rho^{4+2i} \right). \quad (10)$$

That is absolutely and equally met in circle  $|\rho| < R$  under whole values of parameter  $m$ . To calculate coefficients of the degrees of the sedate line (10) let us build recurrent formulas

$$A_{nm} = \frac{2}{\pi R z_0} \int_{\Gamma} q(\varphi, z) \sin \frac{m\pi z}{z_0} \cos(m\varphi) d\Gamma, \quad n = 1, 2, 3, \dots,$$

$$B_{nm} = \frac{2}{\pi R z_0} \int_{\Gamma} f(\varphi, z) \sin \frac{m\pi z}{z_0} \sin(m\varphi) d\Gamma, \quad n, m = 0, 1, 2, 3, \dots \quad (16)$$

The convergence of the line (15) is proved with a principle of maximum for elliptic equations.

### Multiple polynomials

To boost the process of calculating coefficients of  $P_{ij}(m)$  line (10) let us introduce auxiliary functions [5]

$$Q_{i,l} = \sum_{\tau_1=0}^{i-2-2l} \sum_{\tau_2=0}^{\tau_1} \dots \sum_{\tau_{l+1}=0}^{\tau_l} \prod_{\eta=0}^l (2 + 2\eta + \tau_{l+1-\eta}). \quad (17)$$

That represent simplified modification of the multiple polynomials (13).

Then for whole nonnegative values  $l \geq 1$ ,  $i > 2 + 2l$ ,  $\tau_i$  sequence of multipliers into sum-

mands of function  $Q_{i,l}$  with similar values of  $l$  can be written as block matrixes of triangle shape. For example, while  $l = 1$ ,  $i = 5, 6, 7$  и  $l = 2$ ,  $i = 6, 7, 8$  we have summs

$$Q_{5,1} = 2 \cdot 4 + 2 \cdot 5 + 3 \cdot 5, \quad Q_{6,1} = 2 \cdot 4 + 2 \cdot 5 + 3 \cdot 5 + 2 \cdot 6 + 3 \cdot 6 + 4 \cdot 6,$$

$$Q_{7,1} = 2 \cdot 4 + 2 \cdot 5 + 3 \cdot 5 + 2 \cdot 6 + 3 \cdot 6 + 4 \cdot 6 + 2 \cdot 7 + 3 \cdot 7 + 4 \cdot 7 + 5 \cdot 7,$$

$$Q_{7,2} = 2 \cdot 4 \cdot 6 + 2 \cdot 4 \cdot 7 + 2 \cdot 5 \cdot 7 + 3 \cdot 5 \cdot 7,$$

$$Q_{8,2} = 2 \cdot 4 \cdot 6 + 2 \cdot 4 \cdot 7 + 2 \cdot 5 \cdot 7 + 3 \cdot 5 \cdot 7 + 2 \cdot 4 \cdot 8 + 2 \cdot 5 \cdot 8 + 3 \cdot 5 \cdot 8 + 2 \cdot 6 \cdot 8 + 3 \cdot 6 \cdot 8 + 4 \cdot 6 \cdot 8,$$

to which will correspond the matrixes

$$Q_{5,1} \rightarrow \begin{pmatrix} 2 \cdot 4 & 2 \cdot 5 \\ 0 & 3 \cdot 5 \end{pmatrix}; \quad Q_{6,1} \rightarrow \begin{pmatrix} 2 \cdot 4 & 2 \cdot 5 & 2 \cdot 6 \\ 0 & 3 \cdot 5 & 3 \cdot 6 \\ 0 & 0 & 4 \cdot 6 \end{pmatrix};$$

$$Q_{7,1} \rightarrow \begin{pmatrix} 2 \cdot 4 & 2 \cdot 5 & 2 \cdot 6 & 2 \cdot 7 \\ 0 & 3 \cdot 5 & 3 \cdot 6 & 3 \cdot 7 \\ 0 & 0 & 4 \cdot 6 & 4 \cdot 7 \\ 0 & 0 & 0 & 5 \cdot 7 \end{pmatrix}; \quad Q_{7,2} \rightarrow \begin{pmatrix} 2 \cdot 4 \cdot 6 & 2 \cdot 4 \cdot 7 & 3 \cdot 5 \cdot 7 \\ 0 & 2 \cdot 5 \cdot 7 & 0 \end{pmatrix};$$

$$Q_{8,2} \rightarrow \begin{pmatrix} 2 \cdot 4 \cdot 6 & 2 \cdot 4 \cdot 7 & 2 \cdot 4 \cdot 8 & 3 \cdot 5 \cdot 7 & 3 \cdot 5 \cdot 8 & 4 \cdot 6 \cdot 8 \\ 0 & 2 \cdot 5 \cdot 7 & 2 \cdot 5 \cdot 8 & 0 & 3 \cdot 6 \cdot 8 & 0 \\ 0 & 0 & 2 \cdot 6 \cdot 8 & 0 & 0 & 0 \end{pmatrix}.$$

In connection with this characteristic functions  $P_{i,l}(n)$  are called *multiple multinomials of triangle form*.

### Resume

While modeling physical processes in extreme conditions the most basic and difficult stage is the correct setting of an objective. In this work we have obtained the terms that provide for existence and uniqueness of the solution of the first boundary for degenerating on the line elliptic equations. During the problem research a special function class has been built and called multiple polynomials of triangle form.

### References

1. Nesmayanov A.A., Sigalov G.F. Diffraction of small-water waves on thin body under near-critical movement speed // Young scientists' messenger. / Series: applied maths and mechanics. –SPb: Ed. SPB state technical university, 2002, №1. – P. 72-77.

2. Sergiyenko L.S. On setting of correct objectives for degenerating models of stationary processes that take place in solenoid speed field // Non-classic equation of mathematic physics. – Novosibirsk: Ed. Institute of mathematics, 2002. – P. 226-230.

3. Sergiyenko L.S. Mathematic modeling of physical-technical processes. – Irkutsk: Ed. Irkutsk state technical university, 2006. – P. 58-91, 169-173.

4. Sergiyenko L.S. On problem of Dirichlet for one class of degenerating on axis elliptic equations // Modern methods of function theory and adjoining problems: materials of Voronezh winter mathematic school / Voronezh state university, Moscow state university of M.V. Lomonosov, Mathematic institute of V.A. Steklov of Russian science academy: Voronezh: Ed. centre of Voronezh state university. – 2011. – P. 311-312

5. Sergiyenko L.S., Byenchayeva A.V. The first boundary problem for stationary equation of Schrodinger class // Irkutsk state technical university Messenger / scientific magazine – Irkutsk: Ed. Irkutsk state technical university, 2011. – № 10. – P. 275-280.

6. Tichonov A.N., Samarskiy A.A. Equations of mathematic physics: textbook. –M.: Science, 1977. –P. 484-495, 518-527, 615-655.

7. Janushauskas A.I. Analytic theory of elliptic equations. – Novosibirsk: Science, 1978. – P. 159-162.